

Westminster House, 49 Knowsley Street, Bury, Greater Manchester, BL9 0ST.
Tel: 0161-762-9108 Fax: 0161-272-0292
www.medical-specialists.co.uk

Please read this leaflet before you use your medicine. This leaflet provides some useful information for you on your medicine. If you have any questions, please contact us. No liability is accepted for misinterpretation of this information.

What is Xifaxanta (Rifaximin)

It is an anti-biotic that stays and acts locally in the intestine, and is not absorbed into the blood stream.

It is licensed for the use in Travellers' Diarrhoea in the UK but not for Irritable Bowel Syndrome (IBS), although in America, the Federal Drug Administration (FDA) have approved it for the treatment of IBS.

We strongly advise you to read the enclosed Patient Information Leaflet and read the Contra-indications and possible Side-effects, and if you have any of these, stop the medication immediately and contact us.

How to take Xifaxanta for IBS/SIBO

Take TWO tablets THREE times a day, OR As Directed.

We also strongly recommend you go back to the product page for Xifaxanta on our website and click on the link contained in the paragraph about IBS, which takes you to information and dosage recommended for IBS by Dr Mark Hyman. He also gives '**5 Steps to Curing IBS**' information. He also advises about getting an IgG blood test, diet probiotics and much more.

Westminster House, 49 Knowsley Street, Bury, Greater Manchester, BL9 0ST.
Tel: 0161-762-9108 Fax: 0161-272-0292
www.medical-specialists.co.uk

How to take Xifaxanta for IBS/SIBO (High-Strength Courses)

10 days:

Take THREE tablets THREE times a day for 10 days, OR As Directed.

14 days:

Take THREE tablets THREE times a day for 14 days, OR As Directed.